


## Atoms, Molecules and Ions

Chapter 2



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


## Dalton's Atomic Theory (1808)

1. Elements are composed of extremely small particles called **atoms**.
2. All **atoms** of a given element are identical, having the same size, mass and chemical properties. The atoms of one element are different from the atoms of all other elements.
3. **Compounds** are composed of atoms of more than one element. In any compound, the ratio of the numbers of atoms of any two of the elements present is either an integer or a simple fraction.
4. A **chemical reaction** involves only the separation, combination, or rearrangement of atoms; it does not result in their creation or destruction.


2

## Dalton's Atomic Theory

Carbon monoxide




Carbon dioxide




Law of Multiple Proportions


3



Atoms of element X



Atoms of element Y




Compounds of elements X and Y



Law of Conservation of Mass

4

## Cathode Ray Tube



J.J. Thomson, **measured mass/charge of  $e^-$**   
(1906 Nobel Prize in Physics) 5

## Cathode Ray Tube



6

### Millikan's Experiment

**Measured mass of  $e^-$**   
(1923 Nobel Prize in Physics)

$e^-$  charge =  $-1.60 \times 10^{-19}$  C  
 Thomson's charge/mass of  $e^-$  =  $-1.76 \times 10^8$  C/g  
 $e^-$  mass =  $9.10 \times 10^{-28}$  g

### Types of Radioactivity

### Thomson's Model

### Rutherford's Experiment

(1908 Nobel Prize in Chemistry)

$\alpha$  particle velocity  $\sim 1.4 \times 10^7$  m/s  
( $\sim 5\%$  speed of light)

- atoms positive charge is concentrated in the nucleus
- proton (p) has opposite (+) charge of electron (-)
- mass of p is 1840 x mass of  $e^-$  ( $1.67 \times 10^{-24}$  g)

### Rutherford's Model of the Atom

atomic radius  $\sim 100$  pm =  $1 \times 10^{-10}$  m  
 nuclear radius  $\sim 5 \times 10^{-3}$  pm =  $5 \times 10^{-15}$  m

"If the atom is the Houston Astrodome, then the nucleus is a marble on the 50-yard line."

### Chadwick's Experiment (1932)

(1935 Noble Prize in Physics)

H atoms - 1 p; He atoms - 2 p  
 mass He/mass H should = 2  
 measured mass He/mass H = 4

$\alpha + {}^9\text{Be} \longrightarrow {}^1_0\text{n} + {}^{12}\text{C} + \text{energy}$

neutron (n) is neutral (charge = 0)  
 n mass  $\sim$  p mass =  $1.67 \times 10^{-24}$  g

**TABLE 2.1** Mass and Charge of Subatomic Particles

Particle	Mass (g)	Charge	
		Coulomb	Charge Unit
Electron <sup>e-</sup>	$9.10938 \times 10^{-28}$	$-1.6022 \times 10^{-19}$	-1
Proton	$1.67262 \times 10^{-24}$	$+1.6022 \times 10^{-19}$	+1
Neutron	$1.67493 \times 10^{-24}$	0	0

More refined measurements have given us a more accurate value of an electron's mass than Millikan's.

mass p    mass n    1840 x mass e<sup>-</sup>

13

**Atomic number, Mass number and Isotopes**

**Atomic number (Z)** = number of protons in nucleus

**Mass number (A)** = number of protons + number of neutrons  
= atomic number (Z) + number of neutrons


**Isotopes** are atoms of the same element (X) with different numbers of neutrons in their nuclei

Mass Number → A X ← Element Symbol  
Atomic Number → Z X ← Element Symbol

${}^1_1\text{H}$      ${}^2_1\text{H}$  (D)     ${}^3_1\text{H}$  (T)

${}^{235}_{92}\text{U}$      ${}^{238}_{92}\text{U}$

14



15

How many protons, neutrons, and electrons are in  ${}^{14}_6\text{C}$ ?

6 protons, 8 (14 - 6) neutrons, 6 electrons

How many protons, neutrons, and electrons are in  ${}^{11}_6\text{C}$ ?

6 protons, 5 (11 - 6) neutrons, 6 electrons


16

**The Modern Periodic Table**

Period


Metals    Metalloids    Nonmetals

17




18

A **molecule** is an aggregate of two or more atoms in a definite arrangement held together by chemical forces



A **diatomic molecule** contains only two atoms.

H<sub>2</sub>, N<sub>2</sub>, O<sub>2</sub>, Br<sub>2</sub>, HCl, CO



diatomic elements

A **polyatomic molecule** contains more than two atoms

O<sub>3</sub>, H<sub>2</sub>O, NH<sub>3</sub>, CH<sub>4</sub>

19

An **ion** is an atom, or group of atoms, that has a net positive or negative charge.

**cation** – ion with a positive charge

If a neutral atom **loses** one or more electrons it becomes a cation.



11 protons  
11 electrons



11 protons  
10 electrons

**anion** – ion with a negative charge

If a neutral atom **gains** one or more electrons it becomes an anion.



17 protons  
17 electrons



17 protons  
18 electrons

A **monatomic ion** contains only one atom


Na<sup>+</sup>, Cl<sup>-</sup>, Ca<sup>2+</sup>, O<sup>2-</sup>, Al<sup>3+</sup>, N<sup>3-</sup>

A **polyatomic ion** contains more than one atom

OH<sup>-</sup>, CN<sup>-</sup>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>

21

### Common Ions Shown on the Periodic Table



1A	2A	3A	4A	5A	6A	7A	8A	9A	10A	11A	12A	13A	14A	15A	16A	17A	18A
Li <sup>+</sup>												Al <sup>3+</sup>		P <sup>3-</sup>	S <sup>2-</sup>	Cl <sup>-</sup>	
K <sup>+</sup>	Ca <sup>2+</sup>				Cy <sup>2+</sup> Cy <sup>3+</sup>	Mn <sup>2+</sup> Mn <sup>3+</sup>	Fe <sup>2+</sup> Fe <sup>3+</sup>	Cu <sup>2+</sup> Cu <sup>+</sup>	Ni <sup>2+</sup> Ni <sup>3+</sup>	Cd Cu <sup>+</sup>	Zn <sup>2+</sup>			Se <sup>2-</sup> Se <sup>3-</sup>	Te <sup>2-</sup>	Br <sup>-</sup>	
Rb <sup>+</sup>	Sr <sup>2+</sup>									Ag <sup>+</sup>	Cd <sup>2+</sup>			Sn <sup>2+</sup> Sn <sup>4+</sup>		I <sup>-</sup>	
Cs <sup>+</sup>	Ba <sup>2+</sup>									Au <sup>+</sup> Au <sup>3+</sup>	Hg <sup>2+</sup> Hg <sup>1+</sup>			Pb <sup>2+</sup> Pb <sup>4+</sup>			

22

How many protons and electrons are in  ${}_{13}^{27}\text{Al}^{3+}$  ?

13 protons, 10 (13 – 3) electrons

How many protons and electrons are in  ${}_{34}^{78}\text{Se}^{2-}$  ?

34 protons, 36 (34 + 2) electrons

23

### Formulas and Models

	Hydrogen	Water	Ammonia	Methane
Molecular formula	H <sub>2</sub>	H <sub>2</sub> O	NH <sub>3</sub>	CH <sub>4</sub>
Structural formula	H—H	H—O—H	H—N—H   H	H—C—H   H
Ball-and-stick model				
Space-filling model				

24

A **molecular formula** shows the exact number of atoms of each element in the smallest unit of a substance

An **empirical formula** shows the simplest whole-number ratio of the atoms in a substance



<u>molecular</u>	<u>empirical</u>
H <sub>2</sub> O	H <sub>2</sub> O
C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	CH <sub>2</sub> O
O <sub>3</sub>	O
N <sub>2</sub> H <sub>4</sub>	NH <sub>2</sub>

25

**ionic compounds** consist of a combination of cations and anions

- The formula is usually the same as the empirical formula
- The sum of the charges on the cation(s) and anion(s) in each formula unit must equal zero

The ionic compound NaCl






26

1A	2A	3A	4A	5A	6A	7A	8A
Li				N	O	F	
Na	Mg	Al		S	Cl		
K	Ca					Br	
Rb	Sr					I	
Cs	Ba						

The most reactive **metals** (green) and the most reactive **nonmetals** (blue) combine to form ionic compounds.

27

### Formula of Ionic Compounds



28

### Chemical Nomenclature

#### • Ionic Compounds

- Often a metal + nonmetal
- Anion (nonmetal), add “ide” to element name

BaCl <sub>2</sub>	barium chloride
K <sub>2</sub> O	potassium oxide
Mg(OH) <sub>2</sub>	magnesium hydroxide
KNO <sub>3</sub>	potassium nitrate

29

- Transition metal ionic compounds
  - indicate charge on metal with Roman numerals

FeCl <sub>2</sub>	2 Cl <sup>-</sup> -2 so Fe is +2	iron(II) chloride
FeCl <sub>3</sub>	3 Cl <sup>-</sup> -3 so Fe is +3	iron(III) chloride
Cr <sub>2</sub> S <sub>3</sub>	3 S <sup>-2</sup> -6 so Cr is +3 (6/2)	chromium(III) sulfide

30

**TABLE 2.2** The “-ide” Nomenclature of Some Common Monatomic Anions According to Their Positions in the Periodic Table

Group 4A	Group 5A	Group 6A	Group 7A
C carbide ( $C^{4-}$ ) <sup>a</sup>	N nitride ( $N^{3-}$ )	O oxide ( $O^{2-}$ )	F fluoride ( $F^{-}$ )
Si silicide ( $Si^{4-}$ )	P phosphide ( $P^{3-}$ )	S sulfide ( $S^{2-}$ )	Cl chloride ( $Cl^{-}$ )
		Se selenide ( $Se^{2-}$ )	Br bromide ( $Br^{-}$ )
		Te telluride ( $Te^{2-}$ )	I iodide ( $I^{-}$ )

<sup>a</sup>The word “carbide” is also used for the anion  $C_2^{2-}$ .

**TABLE 2.3** Names and Formulas of Some Common Inorganic Cations and Anions

Cation	Anion
ammonium ( $NH_4^+$ )	bicarbonate ( $HCO_3^-$ )
ammonium hydroxide ( $NH_4OH$ )	carbonate ( $CO_3^{2-}$ )
barium ( $Ba^{2+}$ )	chlorate ( $ClO_3^-$ )
bromine ( $Br_2^{+}$ )	chloride ( $Cl^-$ )
calcium ( $Ca^{2+}$ )	chromate ( $CrO_4^{2-}$ )
cesium ( $Cs^+$ )	cyanide ( $CN^-$ )
dibromine ( $Br_2^{2+}$ )	dichromate ( $Cr_2O_7^{2-}$ )
dichlorine ( $Cl_2^{2+}$ )	diphosphate ( $P_2O_7^{4-}$ )
diphosphorus ( $P_2^{2+}$ )	diphosphite ( $P_2O_5$ )
dipropyl ( $C_6H_{14}^{2+}$ )	hydrogen carbonate or bicarbonate ( $HCO_3^-$ )
hydrogen ( $H^+$ )	hydrogen phosphate ( $H_2PO_4^-$ )
iron(II) or ferrous ( $Fe^{2+}$ )	hydrogen phosphite ( $H_2PO_3^-$ )
iron(III) or ferric ( $Fe^{3+}$ )	hydrogen sulfite or bisulfite ( $HSO_3^-$ )
lead(II) or plumbous ( $Pb^{2+}$ )	hydroxide ( $OH^-$ )
lead(IV) or plumbic ( $Pb^{4+}$ )	iodide ( $I^-$ )
lithium ( $Li^+$ )	iodate ( $IO_3^-$ )
magnesium ( $Mg^{2+}$ )	iodate ( $IO_4^-$ )
magnesium(II) or magnesian ( $Mg^{2+}$ )	iodide ( $I^-$ )
manganese(II) or manganous ( $Mn^{2+}$ )	iodine ( $I_2$ )
manganese(IV) or manganic ( $Mn^{4+}$ )	iodine ( $I_2$ )
mercury(II) or mercurous ( $Hg_2^{2+}$ )	iodine ( $I_2$ )
mercury(IV) or mercuric ( $Hg^{2+}$ )	iodine ( $I_2$ )
potassium ( $K^+$ )	isocyanate ( $NCN^-$ )
radium ( $Ra^{2+}$ )	nitrate ( $NO_3^-$ )
silver ( $Ag^+$ )	nitrite ( $NO_2^-$ )
sodium ( $Na^+$ )	nitrite ( $NO_2^-$ )
strontium ( $Sr^{2+}$ )	nitrite ( $NO_2^-$ )
tin(II) or stannous ( $Sn^{2+}$ )	nitrite ( $NO_2^-$ )
tin(IV) or stannic ( $Sn^{4+}$ )	nitrite ( $NO_2^-$ )
zinc ( $Zn^{2+}$ )	nitrite ( $NO_2^-$ )

**Molecular compounds**


- Nonmetals or nonmetals + metalloids
- Common names
  - $H_2O$ ,  $NH_3$ ,  $CH_4$ ,
- Element furthest to the left in a period and closest to the bottom of a group on periodic table is placed first in formula
- If more than one compound can be formed from the same elements, use prefixes to indicate number of each kind of atom
- Last element name ends in *ide*

**TABLE 2.4** Greek Prefixes Used in Naming Molecular Compounds

Prefix	Meaning
mono-	1
di-	2
tri-	3
tetra-	4
penta-	5
hexa-	6
hepta-	7
octa-	8
nona-	9
deca-	10

**Molecular Compounds**

HI	hydrogen iodide
NF <sub>3</sub>	nitrogen trifluoride
SO <sub>2</sub>	sulfur dioxide
N <sub>2</sub> Cl <sub>4</sub>	dinitrogen tetrachloride
NO <sub>2</sub>	nitrogen dioxide
N <sub>2</sub> O	dinitrogen monoxide



An **acid** can be defined as a substance that yields hydrogen ions ( $H^+$ ) when dissolved in water.

For example:  $HCl$  gas and  $HCl$  in water


- Pure substance, hydrogen chloride  $HCl$
- Dissolved in water ( $H_3O^+$  and  $Cl^-$ ), hydrochloric acid

TABLE 2.5 Some Simple Acids	
Anion	Corresponding Acid
F <sup>-</sup> (fluoride)	HF (hydrofluoric acid)
Cl <sup>-</sup> (chloride)	HCl (hydrochloric acid)
Br <sup>-</sup> (bromide)	HBr (hydrobromic acid)
I <sup>-</sup> (iodide)	HI (hydroiodic acid)
CN <sup>-</sup> (cyanide)	HCN (hydrocyanic acid)
S <sup>2-</sup> (sulfide)	H <sub>2</sub> S (hydrosulfuric acid)


37

An **oxoacid** is an acid that contains hydrogen, oxygen, and another element.

HNO<sub>3</sub> nitric acid



H<sub>2</sub>CO<sub>3</sub> carbonic acid




H<sub>3</sub>PO<sub>4</sub> phosphoric acid



38

### Naming Oxoacids and Oxoanions



39

The rules for naming **oxoanions**, *anions of oxoacids*, are as follows:

1. When all the H ions are removed from the "-ic" acid, the anion's name ends with "-ate."
2. When all the H ions are removed from the "-ous" acid, the anion's name ends with "-ite."
3. The names of anions in which one or more but not all the hydrogen ions have been removed must indicate the number of H ions present.

For example:

- H<sub>2</sub>PO<sub>4</sub><sup>-</sup> dihydrogen phosphate
- HPO<sub>4</sub><sup>2-</sup> hydrogen phosphate
- PO<sub>4</sub><sup>3-</sup> phosphate

40

TABLE 2.6 Names of Oxoacids and Oxoanions That Contain Chlorine	
Acid	Anion
HClO <sub>4</sub> (perchloric acid)	ClO <sub>4</sub> <sup>-</sup> (perchlorate)
HClO <sub>3</sub> (chloric acid)	ClO <sub>3</sub> <sup>-</sup> (chlorate)
HClO <sub>2</sub> (chlorous acid)	ClO <sub>2</sub> <sup>-</sup> (chlorite)
HClO (hypochlorous acid)	ClO <sup>-</sup> (hypochlorite)

41

A **base** can be defined as a substance that yields hydroxide ions (OH<sup>-</sup>) when dissolved in water.

NaOH sodium hydroxide

KOH potassium hydroxide

Ba(OH)<sub>2</sub> barium hydroxide

42

**Hydrates** are compounds that have a specific number of water molecules attached to them.

$\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$  barium chloride dihydrate

$\text{LiCl} \cdot \text{H}_2\text{O}$  lithium chloride monohydrate

$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  magnesium sulfate heptahydrate

$\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  strontium nitrate tetrahydrate

$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  →  ←  $\text{CuSO}_4$

43


TABLE 2.7 Common and Systematic Names of Some Compounds

Formula	Common Name	Systematic Name
$\text{H}_2\text{O}$	Water	Dihydrogen monoxide
$\text{NH}_3$	Ammonia	Trihydrogen nitride
$\text{CO}_2$	Dry ice	Solid carbon dioxide
$\text{NaCl}$	Table salt	Sodium chloride
$\text{N}_2\text{O}$	Laughing gas	Dinitrogen monoxide
$\text{CaCO}_3$	Marble, chalk, limestone	Calcium carbonate
$\text{CaO}$	Quicklime	Calcium oxide
$\text{Ca}(\text{OH})_2$	Slaked lime	Calcium hydroxide
$\text{NaHCO}_3$	Baking soda	Sodium hydrogen carbonate
$\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$	Washing soda	Sodium carbonate decahydrate
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	Epsom salt	Magnesium sulfate heptahydrate
$\text{Mg}(\text{OH})_2$	Milk of magnesia	Magnesium hydroxide
$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Gypsum	Calcium sulfate dihydrate

44

**Organic chemistry** is the branch of chemistry that deals with carbon compounds


### Functional Groups



methanol

methylamine

acetic acid



$\text{CH}_3\text{OH}$



$\text{CH}_3\text{NH}_2$



$\text{CH}_3\text{COOH}$

45

TABLE 2.8 The First Ten Straight-Chain Alkanes

Name	Formula	Molecular Model
Methane	$\text{CH}_4$	
Ethane	$\text{C}_2\text{H}_6$	
Propane	$\text{C}_3\text{H}_8$	
Butane	$\text{C}_4\text{H}_{10}$	
Pentane	$\text{C}_5\text{H}_{12}$	
Hexane	$\text{C}_6\text{H}_{14}$	
Heptane	$\text{C}_7\text{H}_{16}$	
Octane	$\text{C}_8\text{H}_{18}$	
Nonane	$\text{C}_9\text{H}_{20}$	
Decane	$\text{C}_{10}\text{H}_{22}$	

46